



North Pacific Fisheries Commission

Species summary for Japanese sardine

Japanese sardine (*Sardinops melanostictus*)

Common names:

拟沙丁鱼, Ni Sha Ding Yu (China)

マイワシ, Maiwashi (Japan)

정어리, Jeong-eoli (Korea)

Дальневосточная сардина (Russia)

遠東擬沙丁魚, Yuan-Dong-Ni-Sha-Ding-Yu (Chinese Taipei)



Figure 1. Japanese Sardine (*Sardinops melanostictus*).

Management

Active NPFC Management Measures

The following NPFC conservation and management measure (CMM) pertains to this species:

- CMM 2024-11 For Japanese Sardine, Neon Flying Squid and Japanese Flying Squid

Available from <https://www.npfc.int/active-conservation-and-management-measures>

Management Summary

The current management measure for Japanese Sardine does not specify catch or effort limits. The CMM states that Members and Cooperating non-Contracting Parties currently harvesting Japanese Sardine should refrain from expansion of the number of fishing vessels authorized to fish Japanese Sardine in the Convention Area. New harvest capacity should also be avoided until as stock assessment has been completed.

A stock assessment for Japanese Sardine is conducted by Japan within their EEZ and used for management of the domestic fishery.


Table 1. Current status of NPFC management measures

Convention or Management Principle	Status	Comment or Consideration
Biological reference point(s)	Not accomplished	Not established for NPFC CA (Established in Japan EEZ)
Stock status	Unknown	Status determination criteria not established for NPFC CA (Established in Japan EEZ)
Catch limit	Intermediate	Not established for NPFC CA (Recommended catch, effort limits in Japan EEZ)
Harvest control rule	Not accomplished	Not established for NPFC CA (Established in Japan EEZ)
Other	Intermediate	No expansion of fishing beyond currently fished areas

Assessment

There is currently no stock assessment for Japanese Sardine conducted by NPFC for the Convention Area.

Japan conducts an assessment of the Japanese Sardine stock using ridge VPA and a number of data sources described below (Hiroshi and Nishida 2005). The catch and biomass estimated by Japan's stock assessment have both been increasing since 2010 (Figure 3). The spawning stock biomass is currently estimated to be higher than SSB_{msy} , but fishing mortality is higher than F_{msy} indicating overfishing in the most recent 3 years (Figure 3). Japan's stock domestic assessment includes catch data from the NPFC CA by China and Russia. Information on the size, weight and age of the catch from the NPFC CA would be useful if it were made available for Japan's stock assessment.






Figure 3. Time series of catch by age in the Japanese Sardine fishery and time series of spawning stock biomass, total biomass and exploitation rate from the domestic Japanese Sardine stock assessment (top panels). Kobe plot indicating historical and current status of Japanese sardine in relation to MSY-based reference points (reprinted from Japan's domestic stock assessment of Japanese Sardine (bottom panel)).

Data

Surveys

Japan conducts three surveys that estimate recruitment for a number of pelagic species, including Japanese Sardine (Table 2). The surveys target pre-recruits and juveniles to determine an index of recruitment. Japan also conducts a monthly egg and larval survey that is used to estimate spawning stock biomass. Surveys are conducted in spring (1995-2024), summer (2001-2024) and fall (2005-2024) at 30-80 stations per year. The survey protocol can be found at (Oozeki et al. 2007). Russia has conducted a summertime acoustic-trawl survey since 2010 that examines mid-water and upper epipelagic species including Japanese Sardine. China has been conducting a scientific survey using its fishery research vessel Song Hang in the convention area of NPFC since 2021. The survey is conducted during June-August, with methods of mid-trawling, acoustic and squid jigging, covering about 70 stations per year.

Fishery

China, Japan and Russia catch Japanese sardine. China does not target the species, but it is captured as bycatch in other fisheries (e.g. chub mackerel). Catches are primarily by purse seine, with a smaller component of the catch taken by pelagic trawl. China's catch of Japanese Sardine is taken exclusively from the Convention Area from April to December. China's existing catch records are from 2016 to 2024 and show increasing catches during that time period as the stock

may have been increasing. The historical catches (prior to 2016) are unknown, thought to be low and likely need to be confirmed.

Japan's fishery for Japanese Sardine occurs inside their EEZ and is mostly conducted by large purse seine vessels (>90% of the catch). Additional components of the fishery include set nets, dip nets and other gears. The fishery experienced very high catches in the 1980's and early 1990's, a decline to very low catches from 1995 to ~2010 and has been recovering since then. The fishery is conducted year round, but mainly during the summer season.

The Russian fishery occurs inside their EEZ and is prosecuted primarily by pelagic trawling (>90% of the catch), with a smaller component of the catch coming from purse seines. The success of Russian fishery depends on the migration patterns and overall abundance of Japanese Sardine, as the sardine move into Russian waters when their abundance is high. For this reason, there was no catch from 1994-2011 when the stock abundance was low, but in recent years (since 2016) as the stock has recovered and water temperatures have been warm there have been increasing catches in Russia. The Russian fishery occurs primarily from June to November.




Figure 4. Historical catch of Japanese Sardine by Members in both the CA and inside Members EEZs.

Other NPFC Members (Canada, EU, Korea, Chinese Taipei, USA and Vanuatu) do not target Japanese Sardine. Chinese Taipei has some historical records of Japanese Sardine bycatch in the Pacific Saury fishery (~100 mt) and Korea has a small amount of historical bycatch data from the bottom trawl fishery. Vanuatu, USA, EU and Canada have no record of Japanese Sardine catches. However, there have been recent (since 2022) occurrences of Japanese Sardine along the USA west coast.

Fishery catch data is available for Members from the NPFC website (<https://www.npfc.int/system/files/2024-04/NPFC-2024-AR-Annual%20Summary%20Footprint%20-%20Japanese%20Sardine.xlsx>) since 2001. Prior years fishery catch data was downloaded from FAO data collections at <https://www.openfisheries.org> using rfisheries package (Karthik Ram, Carl Boettiger, and Dyck 2013).




Figure 5. Historical fishing effort for Japanese Sardine.

Biological collections

China has collected biological data from fishery catches of Japanese Sardine since 2020. These collections included length data as well as maturity and age structures.

Russia collects length and weight data, age structures (scales) and maturity data from both commercial catches and surveys.

Japan also collects length, weight, maturity and age data from the survey and fishery to support their stock assessment.

Data availability from Members regarding Japanese sardine

Data	Source	Years	Comment
Catch	China	2015-present	Catches from convention area
	Japan	1995-present	Historical catch data from 1968 available, catches in national waters
	Korea		Minor bycatch in bottom trawl fishery
	Russia	2016-present	Catches primarily in national waters, not convention area
	Chinese		Minor bycatch in Pacific saury fishery

Data	Source	Years	Comment
	Taipei		
CPUE			not developed
Survey	Japan		Pre-recruit survey
	Japan		Juvenile survey
	Japan		Monthly egg and larval survey
	Russia	2010-present	Acoustic-trawl survey
	China	2021-present	Midwater trawling and acoustics
Age data	China	2020-present	Commercial catch
	Japan		Commercial and survey catches
	Russia		Commercial and survey catches
Length data	China	2020-present	Commercial catch
	Japan		Commercial and survey catches
	Russia		Commercial and survey catches
Maturity/fecundity	China	2020-present	Commercial catch
	Japan		Commercial and survey catches
	Russia		Commercial and survey catches

Special Comments

None

Biological Information

Distribution

Japanese sardine (*Sardinops melanostictus*; Figure 1) are a pelagic species that occurs in large migratory schools in the coastal waters of China, Chinese Taipei, Japan, Korea and Russia (Figure 4, (Kaschner et al. 2019)). They generally migrate from the south to the north during summer, returning to inshore areas in the south to spawn in the winter. Japanese sardine feed mainly on zooplankton and phytoplankton.

Life history

Japanese sardine are short-lived and fast growing, maturing early at 2-years old. Their maximum length is ~24 cm (Whitehead 1985) and their maximum age is 7 years (Furuichi et al, 2024). Their growth rates and spawning patterns are highly influenced by the environment (Niino et al. 2021) Taxonomically, the Japanese sardine are closely related to other species around the globe including *Sardinops* from southern Africa, Australia, South America and California.




Figure 4. Map of distribution of Sardine species in the North Pacific.

Literature cited

- Hiroshi, and Nishida. 2005. "Stock Assessment and ABC Calculation for Japanese Sardine (*Sardinops Melanostictus*) in the Northwestern Pacific Under Japanese TAC System." In.
- Karthik Ram, Carl Boettiger, and Andrew Dyck. 2013. "Rfisheries: R Interface for Fisheries Data. R Package Version 0.1." 2013. <http://CRAN.R-project.org/package=rfisheries>.
- Kaschner, K., Kesner-Reyes K., Garilao C., Segschneider J., J. Rius-Barile, Rees T., and R. Froese. 2019. "AquaMaps: Predicted Range Maps for Aquatic Species. Data Retrieved from <https://www.aquamaps.org>."
- Furuichi, S., Yukami, R., Kamimura, Y., Nishijima, S., Watanabe, R., Isu, S., & Higashiguchi, K. (2024). Stock assessment and evaluation for Pacific stock of Japanese sardine (fiscal year 2023). Fisheries Agency and Fisheries Research and Education Agency of Japan, Tokyo (2024). https://abchan.fra.go.jp/wpt/wp-content/uploads/2024/03/details_2023_01.pdf
- Niino, Yohei, Sho Furuichi, Yasuhiro Kamimura, and Ryuji Yukami. 2021. "Spatiotemporal spawning patterns and early growth of Japanese sardine in the western North Pacific during the recent stock increase." *Fisheries Oceanography*, no. April: 1–10. <https://doi.org/10.1111/fog.12542>.
- Oozeki, Yoshioki, Akinori Takasuka, Hiroshi Kubota, and Manuel Barange. 2007. "Characterizing Spawning Habitats of Japanese Sardine (*Sardinops Melanostictus*), Japanese Anchovy (*Engraulis Japonicus*), and Pacific Round Herring (*Etrumeus Teres*) in the Northwestern Pacific." *CalCOFI Rep.* 48 (December).
- Whitehead, Peter J. P. 1985. "FAO Species Catalogue. Vol. 7. Clupeoid Fishes of the World (Suborder Clupeoidei). An Annotated and Illustrated Catalogue of the Herrings, Sardines, Pilchards, Sprats, Shads, Anchovies and Wolf-Herrings." *FAO Fish. Synop.* 125(7/1): 1–303.

Appendix: Sardine and the environment

Studies examining the relationship between Japanese sardine and the environment

Reference	Year	Type	Country	Ocean	Region	Species	Life stage	Parameter	Environmental variables	Effect	Method
Kodama, T, Wagawa T, Ohshimo S, Morimoto H, Iguchi N, Fukudome KI, Goto T, Takahashi T, Yasuda T. 2018. Improvement in Recruitment of Japanese Sardine with Delays of the Spring Phytoplankton Bloom in the Sea of Japan. Fisheries Oceanography 27 (4): 289–301. https://doi.org/10.1111/fog.12252 .	2018	journal paper	Japan	Pacific	Sea of Japan	Japanese sardine	Larvae Recruitment	Sea surface chlorophyll a		delay in start and end dates of spring bloom were positively correlated with recruitment	Correlation, n, empirical orthogonal function
Yasuda, Tohya, Satoshi Kitajima, Akira Hayashi, Motomitsu Takahashi, and Masaaki Fukuwaka. 2021. “Cold Offshore Area Provides a Favorable Feeding Ground with Lipid-Rich Foods for Juvenile Japanese Sardine.” Fisheries Oceanography, no. January: 1–16. https://doi.org/10.1111/fog.12530 .	2021	journal paper	Japan	Pacific	Sea of Japan	Japanese juvenile sardine	Body condition	Prey species and temperature		higher condition in offshore distributed fish due to lower temperature and higher lipid content prey	correlation
Nishikawa, Haruka. 2019. “Relationship between Recruitment of Japanese Sardine (Sardinops Melanostictus) and Environment of Larval Habitat in the Low-Stock Period (1995–2010).” Fisheries Oceanography 28 (2): 131–42. https://doi.org/10.1111/fog.12397 .	2019	journal paper	Japan	Pacific	Kuoshi	Japanese sardine	Larvae Recruitment	water temperature and larval drift		warmer temperature related to lower recruitment	correlation
Niino, Yohei, Sho Furuichi, Yasuhiro Kamimura, and Ryuji Yukami. 2021. “Spatiotemporal Spawning Patterns and Early Growth of Japanese Sardine in the Western North Pacific during the Recent Stock Increase.” Fisheries Oceanography, no. April: 1–10. https://doi.org/10.1111/fog.12542 .	2021	journal paper	Japan	Pacific	Kuoshi	Japanese sardine	Larvae growth	spawning distribution and timing (temperature)		early spawning in eastern area contributed to higher recruitment during time of increasing sardine	correlation

Reference	Year	Type	Country	Ocean	Region	Species	Life stage	Parameter	Environmental variables	Effect	Method
Muko, Soyoka, Seiji Ohshimo, Hiroyuki Kurota, Tohya Yasuda, and Masa Aki Fukuwaka. 2018. "Long-Term Change in the Distribution of Japanese Sardine in the Sea of Japan during Population Fluctuations." <i>Marine Ecology Progress Series</i> 593: 141–54. https://doi.org/10.3354/meps12491 .	2018	journal paper	Japan	Pacific	Sea of Japan	Japanese sardine	Adult	Distribution (SDM)	sea surface temperature	biomass dome shaped relationship between sea surface temperature and the probability of presence, with peak between 10-20 C	generalized additive models
Sogawa, Sayaka, Kiyotaka Hidaka, Yasuhiro Kamimura, Masanori Takahashi, Hiroaki Saito, Yuji Okazaki, Yugo Shimizu, and Takashi Setou. 2019. "Environmental Characteristics of Spawning and Nursery Grounds of Japanese Sardine and Mackerels in the Kuroshio and Kuroshio Extension Area." <i>Fisheries Oceanography</i> 28 (4): 454–67. https://doi.org/10.1111/fog.12423 .	2019	journal paper	Japan	Pacific	Kuoshio	Japanese sardine	Egg	Distribution	water temperature, larval drift, zooplankton	little variability in environment where eggs were found, copepod community structure was important	correlation
Kuroda, Hiroshi, Toshihiko Saito, Toshiki Kaga, Akinori Takasuka, Yasuhiro Kamimura, Sho Furuichi, and Takuya Nakanowatari. 2020. "Unconventional Sea Surface Temperature Regime Around Japan in the 2000s–2010s: Potential Influences on Major Fisheries Resources." <i>Frontiers in Marine Science</i> 7 (October): 1–21. https://doi.org/10.3389/fmars.2020.57490 .	2020	journal paper	Japan	Pacific	Japanese	sardine	Adult	Recruitment	PDO, SST	spawning was earlier during SST increases	correlation
Ma, Shuyang, Yongjun Tian, Caihong Fu, Haiqing Yu, Jianchao Li, Yang Liu, Jiahua Cheng, Rong Wan, and Yoshiro	2020	journal paper	China	Pacific	Kuoshio	Japanese sardine	Adult	Abundance/Catch	Basin scale climate (ALPI, SST,	Climate variability introduced	time series analyses

Reference	Year	Type	Country	Ocean	Region	Species	Life stage	Parameter	Environmental variables	Effect	Method
Watanabe. 2021. "Climate-Induced Nonlinearity in Pelagic Communities and Non-Stationary Relationships with Physical Drivers in the Kuroshio Ecosystem." <i>Fish and Fisheries</i> 22 (1): 1–17. https://doi.org/10.1111/faf.12502 .									Current patterns)	nonlinearity and nonstationarity to pelagic fish	
Kurota, Hiroyuki, Cody S. Szuwalski, and Momoko Ichinokawa. 2020. "Drivers of Recruitment Dynamics in Japanese Major Fisheries Resources: Effects of Environmental Conditions and Spawner Abundance." <i>Fisheries Research</i> 221 (September 2019): 105353. https://doi.org/10.1016/j.fishres.2019.105353 .	2020	journal paper	Japan	Pacific	Japanese Pacific	sardine	Adult	Recruitment	"Environment other than SSB	Regime shifts were detected in pelagic species	time series analyses, change point analysis
Furuichi, Sho, Tohya Yasuda, Hiroyuki Kurota, Mari Yoda, Kei Suzuki, Motomitsu Takahashi, and Masa Aki Fukuwaka. 2020. "Disentangling the Effects of Climate and Density-Dependent Factors on Spatiotemporal Dynamics of Japanese Sardine Spawning." <i>Marine Ecology Progress Series</i> 633: 157–68. https://doi.org/10.3354/meps13169 .	2020	journal paper	Japan	Pacific	Sea of Japan	sardine	Egg	Abundance and distribution	SST	Cold water led to decreased egg abundance over larger area, warm temperatures led to earlier spawning	correlation
Okazaki, Yuji, Kazuaki Tadokoro, Hiroshi Kubota, Yasuhiro Kamimura, and Kiyotaka Hidaka. 2019. "Dietary Overlap and Optimal Prey Environments of Larval and Juvenile Sardine and Anchovy in the Mixed Water Region of the Western North Pacific." <i>Marine Ecology Progress Series</i> 630: 149–60. https://doi.org/10.3354/meps13124 .	2019	journal paper	Japan	Pacific	Kuoshio current		larvae and juvenile	prey habits	SST	Temperature influences abundance of prey with effect on recruitment	correlation